Journal of MOLECULAR BIOLOGY

EDITOR-IN-CHIEF
J. C. KENDREW

ASSISTANT EDITOR
S. BRENNER

EDITORIAL BOARD
M. S. Bretscher
C. R. Cantor
P. Chambon
M. Gellert
G. A. Gilbert
M. Gottesman
R. Huber
H. E. Huxley
A. Klug
V. Luzzati
M. F. Moody
P. M. B. Walker
K. Weber
S. M. Weissman

JAN 5 12 12 PM '81
LIBRARY

VOLUME 139
5 May to 5 June 1980

PERIODICAL
QH301
J73

ACADEMIC PRESS
London New York San Francisco

Copyright © 1980, by Academic Press Inc. (London) Ltd.

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 139

Number 1, 5 May 1980

Syvanen, M., Tn903 Induces Inverted Duplications in the Chromosome of Bacteriophage Lambda 1

Robinson, B. H., Lerman, L. S., Beth, A. H., Frisch, H. L., Dalton, L. R. and Auer, C., Analysis of Double-helix Motions with Spin-labeled Probes: Binding Geometry and the Limit of Torsional Elasticity . . . 19

Appendix: Preparation of Spin-labeled Compounds 41

Delbaere, L. T. J. and Brayer, G. D., Structure of the Complex Formed Between the Bacterial-produced Inhibitor Chymostatin and the Serine Enzyme *Streptomyces griseus* Protease A 45

Wright, C. S., Location of the *N*-acetyl-D-neuraminic Acid Binding Site in Wheat Germ Agglutinin. A Crystallographic Study at 2·8 Å Resolution . 53

Minty, A. J. and Gros, F., Coding Potential of Non-polyadenylated Messenger RNA in Mouse Friend Cells 61

Letters to the Editor

Francesco, de L. and Attardi, G., Analysis of Sequence Homology between Human and Mouse Mitochondrial DNA 85

McKay, D. B. and Fried, M. G., Crystallization and Preliminary X-ray Diffraction Data for the Cyclic AMP Receptor Protein of *Escherichia coli* 95

Erratum 97

Number 2, 15 May 1980

Henderson, R. and Shotton, D., Crystallization of Purple Membrane in Three Dimensions 99

Spinelli, G., Melli, M., Arnold, E., Casano, C., Gianguzza, F. and Ciaccio, M., High Molecular Weight RNA Containing Histone Messenger in the Sea Urchin *Paracentrotus lividus* 111

McEwen, B. and Edelstein, S. J., Evidence for a Mixed Lattice in Microtubules Reassembled *in Vitro* 123

Appendix: Crepeau, R. H., Determination of the Degree of Elliptical Flattening for Helical Fibers from Tilted Views 144

Maurer, R., Meyer, B. J. and Ptashne, M., Gene Regulation at the Right Operator (O_R) of Bacteriophage λ. I. O_R3 and Autogenous Negative Control by Repressor 147

MEYER, B. J., MAURER, R. and PTASHNE, M., Gene Regulation at the Right Operator (O_R) of Bacteriophage λ. II. O_R1, O_R2, and O_R3: Their Roles in Mediating the Effects of Repressor and cro 163

Appendix 190

MEYER, B. J. and PTASHNE, M., Gene Regulation at the Right Operator (O_R) of Bacteriophage λ. III. λ Repressor Directly Activates Gene Transcription 195

JANNER, F., VÖGELI, G. and FLURI, R., The Antisuppressor Strain *sin1* of *Schizosaccharomyces pombe* Lacks the Modification Isopentenyladenosine in Transfer RNA 207

KLESSIG, D. F. and CHOW, L. T., Incomplete Splicing and Deficient Accumulation of the Fiber Messenger RNA in Monkey Cells Infected by Human Adenovirus Type 2 221

ISERENTANT, D., VAN MONTAGU, M. and FIERS, W., Studies on the Bacteriophage MS2.XLI. Nature of the Azure Mutation 243

NUMBER 3, 25 MAY 1980

ISHIHARA, A. and HOTANI, H., Micro-video Study of Discontinuous Growth of Bacterial Flagellar Filaments *in Vitro* 265

IMBER, R., TSUGITA, A., WURTZ, M. and HOHN, T., Outer Surface Protein of Bacteriophage Lambda 277

CHARLES, M., SÉMÉRIVA, M. and CHABRE, M., Small-angle Neutron Scattering Study of the Association between Porcine Pancreatic Colipase and Taurodeoxycholate Micelles. 297

PHIZICKY, E. M. and ROBERTS, J. W., Kinetics of *RecA* Protein-directed Inactivation of Repressors of Phage λ and Phage P22 319

RAY, B. K. and APIRION, D., Identification of RNA Molecules which Contain 5 S Ribosomal RNA and Transfer RNA in an RNAase E^- RNAase P^- Double Mutant Strain of *Escherichia coli* 329

GUTHRIE, C. and SCHOLLA, C. A., Asymmetric Maturation of a Dimeric Transfer RNA Precursor 349

FUKADA, K. and ABELSON, J., DNA Sequence of a T4 Transfer RNA Gene Cluster 377

WEBSTER, R. E. and REMENTER, M., Replication of Bacteriophage f1: A Complex Containing Gene *II* Protein in Gene *V* Mutant-infected Bacteria . . 393

LEE, G., HANNETT, N. M., KORMAN, A. and PERO, J., Transcription of Cloned DNA from *Bacillus subtilis* Phage SPO1. Requirement for Hydroxymethyluracil-containing DNA by Phage-modified RNA Polymerase . . . 407

JAMES, M. N. G., BRAYER, G. D., DELBAERE, L. T. J., SIELECKI, A. R. and GERTLER, A., Crystal Structure Studies and Inhibition Kinetics of Tripeptide Chloromethyl Ketone Inhibitors with *Streptomyces griseus* Protease B 423

Bergmann, I. E. and Brawerman, G., Loss of the Polyadenylate Segment from Mammalian Messenger RNA. Selective Cleavage of this Sequence from Polyribosomes 439

Hays, J. B., Korba, B. E. and Konrad, E. B., Novel Mutations of *Escherichia coli* that Produce Recombinogenic Lesions in DNA. I. Identification and Mapping of *arl* Mutations 455

Korba, B. E. and Hays, J. B., Novel Mutants of *Escherichia coli* that Produce Recombinogenic Lesions in DNA. II. Properties of Recombinogenic λ Phages Grown on Bacteria Carrying *arl* Mutations 473

Shick, V. V., Belyavsky, A. V., Bavykin, S. G. and Mirzabekov, A. D., Primary Organization of the Nucleosome Core Particles. Sequential Arrangement of Histones along DNA 491

Belyavsky, A. V., Bavykin, S. G., Goguadze, E. G. and Mirzabekov, A. D., Primary Organization of Nucleosomes Containing all Five Histones and DNA 175 and 165 Base-pairs Long 519

Maquat, L. E., Thornton, K. and Reznikoff, W., *lac* Promoter Mutations Located Downstream from the Transcription Start Site 537

Letters to the Editor

Maquat, L. E. and Reznikoff, W. S., *lac* Promoter Mutation P^r115 Generates a New Transcription Initiation Point 551

Mercolino, T. J., Bellamy, H. D. and Matthews, F. S., A Preliminary Crystallographic Study of β-Glucuronidase from Rat Preputial Gland 557

Mills, A. D., Laskey, R. A., Black, P. and Robertis, E. M. de, An Acidic Protein Which Assembles Nucleosomes *in Vitro* is the Most Abundant Protein in *Xenopus* Oocyte Nuclei. 561

Errata 569

Announcement 571

Number 4, 5 June 1980

Marston, S. B. and Taylor, E. W., Comparison of the Myosin and Actomyosin ATPase Mechanisms of the Four Types of Vertebrate Muscles . . 573

Boyle, J., Robillard, G. T. and Kim, S.-H., Sequential Folding of Transfer RNA. A Nuclear Magnetic Resonance Study of Successively Longer tRNA Fragments with a Common 5′ End 601

Lifson, S. and Sander, C., Specific Recognition in the Tertiary Structure of β-Sheets of Proteins 627

Zackroff, R. V., Weisenberg, R. C. and Deery, J., Equilibrium and Kinetic Analysis of Microtubule Assembly in the Presence of Guanosine Diphosphate 641

Appendix: Weisenberg, R. C., Role of Co-operative Interactions, Microtubule-associated Proteins and Guanosine Triphosphate in Microtubule Assembly: A Model 660

Crawford, J. T. and Goldberg, E. B., The Function of Tail Fibers in Triggering Baseplate Expansion of Bacteriophage T4 679

Jacobs, P., Jauniaux, J.-C. and Grenson, M., A *Cis*-dominant Regulatory Mutation Linked to the *argB-argC* Gene Cluster in *Saccharomyces cerevisiae* 691

Knowlton, R. G., Soll, L. and Yarus, M., Dual Specificity of su^+7 tRNA. Evidence for Translational Discrimination 705

Knowlton, R. G. and Yarus, M., Discrimination Between Aminoacyl Groups on su^+7 tRNA by Elongation Factor Tu 721

Letter to the Editor

Iba, H. and Pkada, Y., Chromosome Segregation in an Asymmetrically Dividing Bacterium, *Caulobacter crescentus* 733

Index to Volume 139

ABELSON, J. (see also FUKUDA, K.), 377
Actomyosin and myosin ATPase mechanisms of the four types of vertebrate muscles, a comparison, 573
Adenovirus type 2-infected monkey cells, incomplete splicing and deficient accumulation of the fiber mRNA, 221
Agglutinin, wheat germ, location of the *N*-acetyl-D-neuraminic acid binding site; crystallographic study at 2.8 Å resolution, 53
Aminoacyl groups on su^+7 tRNA, discrimination between them by elongation factor Tu, 721
Antisuppressor strain *sin1* of *Schizosaccharomyces pombe* lacks the modification isopentenyladenosine in tRNA, 207
APIRION, D. (see also RAY, B. K.), 329
argB-argC gene cluster in *Saccharomyces cerevisiae*, a linked *cis*-dominant regulatory mutation, 691
arl mutations, identification and mapping; novel mutations of *Escherichia coli* that produce recombinogenic lesions in DNA, 455
— — properties of recombinogenic λ phages grown on bacteria carrying these mutations; novel mutants of *Escherichia coli* that produce recombinogenic lesions in DNA, 473
ARNOLD, E. (see also SPINELLI, G.), 111
Assembly of microtubules in the presence of guanosine diphosphate, equilibrium and kinetic analysis, 641
— — —, role of co-operative interactions, microtubule-associated proteins and guanosine triphosphate; a model, 660
ATPase mechanisms of the four types of vertebrate muscles, a comparison, 573
ATTARDI, G. (see also FRANCESCO, DE L.), 85
AUER, C. (see also ROBINSON, B. H.), 19
Azure mutation of bacteriophage MS2, 243

Bacillus subtilis phage SP01, transcription of cloned DNA. Requirement for hydroxymethyluracil-containing DNA by phage-modified RNA polymerase, 407
Bacterial flagellar filaments, micro-video study of discontinuous growth *in vitro*, 265
Bacteriophage F1 replication: a complex containing gene *II* protein in gene *V* mutant-infected bacteria, 393
— λ, gene regulation at the right operator (O_R); λ repressor directly activates gene transcription, 195
— —, — — — — — — — O_R1, O_R2, and O_R3: their roles in mediating the effects of repressor and cro, 163
— —, — — — — — — — (O_R); O_R3 and autogenous negative control by repressor, 147
— — outer surface protein, 277
— — and phage P22, kinetics of *recA* protein-directed inactivation of repressors, 319
— —, recombinogenic, grown on bacteria carrying *arl* mutations; novel mutants of *Escherichia coli* that produce recombinogenic lesions in DNA, 455
— —, Tn903 induces inverted duplications in the chromosome, 1
— MS2, nature of the azure mutation, 243
— SP01, *Bacillus subtilis*, cloned DNA transcription. Requirement for hydroxymethyluracil-containing DNA by phage-modified RNA polymerase, 407
— T4, DNA sequence of a T4 tRNA gene cluster, 377
— —, the function of tail fibers in triggering baseplate expansion, 679
Baseplate expansion of bacteriophage T4, triggering function of tail fibers, 679
BAVYKIN, S. G. (see also BELYAVSKY, A. V.), 519
— — — (see also SHICK, V. V.), 491
BELLAMY, H. D. (see also MERCOLINO, T. J.), 557
BELYAVSKY, A. V., BAVYKIN, S. G., GOGUADZE, E. G., & MIRZABEKOV, A. D., Primary organization of nucleosomes containing all five histones and DNA 175 and 165 base-pairs long, 519
— — — (see also SHICK, V. V.), 491
BERGMANN, I. E. & BRAWERMAN, G., Loss of the polyadenylate segment from mammalian mRNA. Selective cleavage of this sequence from polyribosomes, 439
β-glucuronidase from rat preputial gland, a preliminary crystallographic study, 557
β-sheets of proteins, specific recognition in the tertiary structure, 627

Beth, A. H. (see also Robinson, B. H.), 19
Binding geometry and the limit of torsional elasticity; analysis of double-helix motions with spin-labeled probes, 19
Black, P. (see also Mills, A. D.), 561
Boyle, J., Robillard, G. T. & Kim, S.-H., Sequential folding of transfer RNA. A nuclear magnetic resonance study of successively longer tRNA fragments with a common 5′ end, 601
Brawerman, G. (see also Bergmann, I. E.), 439
Brayer, G.D. (see also Delbaere, L. T. J.), 45
— — — (see also James, M. N. G.), 423

Casano, C. (see also Spinelli, G.), 111
Caulobacter crescentus, chromosome segregation in an asymmetrically dividing bacterium, 733
Chabre, M. (see also Charles, M.), 297
Charles, M., Sémériva, M. & Chabre, M., Small-angle neutron scattering study of the association between porcine pancreatic colipase and taurodeoxycholate micelles, 297
Chow, L. T. (see also Klessig, D. F.), 221
Chromosome of bacteriophage lambda, Tn903 induces inverted duplications, 1
— segregation in an asymmetrically dividing bacterium, *Caulobacter crescentus*, 733
Chymostatin and the serine enzyme *Streptomyces griseus* protease A, structure of the complex formed between them, 45
Ciaccio, M. (see also Spinelli, G.), 111
Cis-dominant regulatory mutation linked to the *argB-argC* gene cluster in *Saccharomyces cerevisiae*, 691
Cloned DNA transcription from *Bacillus subtilis* phage SP01. Requirement for hydroxymethyluracil containing DNA by phage-modified RNA polymerase, 407
Coding potential of non-polyadenylated mRNA in mouse Friend cells, 61
Colipase, porcine pancreatic, small-angle neutron scattering study of the association with taurodeoxycholate micelles, 297
Crawford, J. T. & Goldberg, E. B., The function of tail fibers in triggering baseplate expansion of bacteriophage T4, 679
Crepeau, R. H., Determination of the degree of elliptical flattening for helical fibers from tilted views (*Appendix* to McEwen, B. & Edelstein, S. J., 123), 144
Crystal structure studies and inhibition kinetics of tripeptide chloromethyl ketone inhibitors with *Streptomyces griseus* protease B, 423
Crystallization and preliminary X-ray diffraction data for the cyclic AMP receptor protein of *Escherichia coli*, 95
— of purple membrane in three dimensions, 99
Crystallographic study at 2.8 Å resolution of the location of the *N*-acetyl-D-neuraminic acid binding site in wheat germ agglutinin, 53
— —, preliminary, of β-glucuronidase from rat preputial gland, 557
Cyclic AMP receptor protein of *Escherichia coli*, crystallization and preliminary X-ray diffraction data, 95

Dalton, L. R. (see also Robinson, B. H.), 19
Deery, J. (see also Zackroff, R.V.), 641
Delbaere, L. T. J. & Brayer, G. D., Structure of the complex formed between the bacterial-produced inhibitor chymostatin and the serine enzyme *Streptomyces griseus* protease A, 45
Delbaere, L. T. J. (see also James, M. N. G.), 423
DNA, cloned, transcription from *Bacillus subtilis* phage SP01. Requirement for hydroxymethyluracil-containing DNA by phage-modified RNA polymerase, 407
—, mitochondrial, human and mouse, analysis of sequence homology, 85
—, primary organization of the nucleosome core particles and the sequential arrangement of histones along DNA, 491
— — — — nucleosomes containing all five histones and DNA 175 and 165 base-pairs long, 519
— recombinogenic lesions produced by novel mutations of *Escherichia coli*; identification and mapping of *arl* mutations, 455
— — — — — — — — —; properties of recombinogenic λ phages grown on bacteria carrying *arl* mutations, 473
— sequence of a T4 tRNA gene cluster, 377
Duplications, inverted, induced by Tn903 in the chromosome of bacteriophage lambda, 1

EDELSTEIN, S. J. (see also MCEWEN, B.), 123
Elliptical flattening for helical fibers, determination from tilted views, 144
Elongation factor Tu discriminates between aminoacyl groups on su^+7 tRNA, 721
Escherichia coli, crystallization and preliminary X-ray diffraction data for the cyclic AMP receptor protein, 95
— —, identification of RNA molecules which contain 5 S ribosomal RNA and tRNA in an RNAase E^- RNAase P^- double mutant strain, 329
— —, novel mutants that produce recombinogenic lesions in DNA; properties of recombinogenic λ phages grown on bacteria carrying *arl* mutations, 473
— —, — mutations that produce recombinogenic lesions in DNA; identification and mapping of *arl* mutations, 455

f1 (see also Bacteriophage), 393
Fiber mRNA, incomplete splicing and deficient accumulation in monkey cells infected by human adenovirus type 2, 221
Fibers, helical, determination of the degree of elliptical flattening from tilted views, 144
—, tail, function in triggering baseplate expansion of bacteriophage T4, 679
FIERS, W. (see also ISERENTANT, D.), 243
Filaments, bacterial flagellar, micro-video study of discontinuous growth *in vitro*, 265
Flagellar filaments, bacterial, micro-video study of discontinuous growth *in vitro*, 265
FLURI, R. (see also JANNER, F.), 207
Folding, sequential, of tRNA; n.m.r. study of successively longer tRNA fragments with a common 5′ end, 601
FRANCESCO, DE L. & ATTARDI, G., Analysis of sequence homology between human and mouse mitochondrial DNA, 85
FRIED, M. G. (see also MCKAY, D. B.), 95
Friend cells, mouse, coding potential of non-polyadenylated mRNA, 61
FRISCH, H. L. (see also ROBINSON, B. H.), 19
FUKUDA, K. & ABELSON, J., DNA sequence of a T4 tRNA gene cluster, 377

Gene cluster *argB-argC* in *Saccharomyces cerevisiae*, a linked *cis*-dominant regulatory mutation, 691
— — for T4 tRNA, determination of DNA sequence, 377
— regulation at the right operator (O_R) of bacteriophage λ; λ repressor directly activates gene transcription, 195
— — — — — — — — — — —. O_R1, O_R2, and O_R3: their role in mediating the effects of repressor and cro, 163
— — — — — — — — — — —. O_R3 and autogenous negative control by repressor, 147
— *II* protein contained in complex in gene *V* mutant-infected bacteria; replication of bacteriophage f1, 393
GERTLER, A. (see also JAMES, M. N. G.), 423
GIANGUZZA, F. (see also SPINELLI, G.), 111
GOGUADZE, E. G. (see also BELYAVSKY, A. V.), 519
GOLDBERG, E. B. (see also CRAWFORD, J.T.), 679
GROS, F. (see also MINTY, A. J.), 61
GRENSON, M. (see also JACOBS, P.), 691
Guanosine diphosphate, equilibrium and kinetic analysis of its effect on microtubule assembly, 641
— triphosphate, role in microtubule assembly; a model, 660
GUTHRIE, C. & SCHOLLA, C. A., Asymmetric maturation of a dimeric transfer RNA precursor, 349

HANNETT, N. M. (see also LEE, G.), 407
HAYS, J. B., KORBA, B. E. & KONRAD, E. B., Novel mutations of *Escherichia coli* that produce recombinogenic lesions in DNA. I. Identification and mapping of *arl* mutations, 455
HAYS, J. B. (see also KORBA, B. E.), 473
Helical fibers, determination of the degree of elliptical flattening from tilted views, 144
Helix motions, analysis with spin-labeled probes: binding geometry and the limit of torsional elasticity, 19
HENDERSON, R. & SHOTTON, D., Crystallization of purple membrane in three dimensions, 99

Histone messenger contained in high molecular weight RNA in the sea urchin *Paracentrotus lividus*, 111
— sequential arrangement along DNA; primary organization of the nucleosome core particles, 491
— — — in nucleosomes containing all five histones and DNA 175 and 165 base-pairs long, 519
Hohn, T. (see also Imber, R.), 277
Hotani, H. (see also Ishihara, A.), 265
Human adenovirus type 2-infected monkey cells, incomplete splicing and deficient accumulation of the fiber mRNA, 221
— and mouse mitochondrial DNA, analysis of their sequence homology, 85
Hydroxymethyluracil-containing DNA is required for transcription by phage-modified RNA polymerase; transcription of cloned DNA from *Bacillus subtilis* phage SP01, 407

Iba, H. & Pkada, Y., Chromosome segregation in an asymmetrically dividing bacterium, *Caulobacter crescentus*, 733
Imber, R., Tsugita, A., Wurtz, M. & Hohn, T., Outer surface protein of bacteriophage lambda, 277
Inhibition kinetics and crystal structure studies of tripeptide chloromethyl ketone inhibitors with *Streptomyces griseus* protease B, 423
Iserentant, D., Van Montagu, M. & Fiers, W., Studies on the bacteriophage MS2. XLI. Nature of the azure mutation, 243
Ishihara, A. & Hotani, H., Micro-video study of discontinuous growth of bacterial flagellar filaments *in vitro*, 265
Isopentenyladenosine is missing in tRNA from the antisuppressor strain *sin1* of *Schizosaccharomyces pombe*, 207

Jacobs, P., Jauniaux, J.-C. & Grenson, M., A *cis*-dominant regulatory mutation linked to the *argB-argC* gene cluster in *Saccharomyces cerevisiae*, 691
James, M. N. G., Brayer, G. D., Delbaere, L. T. J., Sielecki, A. R. & Gertler, A., Crystal structure studies and inhibition kinetics of tripeptide chloromethyl ketone inhibitors with *Streptomyces griseus* protease B, 423
Janner, F., Vögeli, G. & Fluri, R., The antisuppressor strain *sin1* of *Schizosaccharomyces pombe* lacks the modification isopentenyladenosine in tRNA, 207
Jauniaux, J.-C. (see also Jacobs, P.), 691

Kim, S.-H. (see also Boyle, J.), 601
Kinetic and equilibrium analysis of microtubule assembly in the presence of guanosine diphosphate, 641
Kinetics of *recA* protein-directed inactivation of repressors of phage λ and phage P22, 319
Klessig, D. F. & Chow, L. T., Incomplete splicing and deficient accumulation of the fiber mRNA in monkey cells infected by human adenovirus type 2, 221
Knowlton, R. G. & Yarus, M., Discrimination between aminoacyl groups on su^+7 tRNA by elongation factor Tu, 721
— — —, Soll, L. & Yarus, M., Dual specificity of su^+7 tRNA. Evidence for translational discrimination, 705
Konrad, E. B. (see also Hays, J. B.), 455
Korba, B. E. & Hays, J. B., Novel mutants of *Escherichia coli* that produce recombinogenic lesions in DNA. II. Properties of recombinogenic λ phages grown on bacteria carrying *arl* mutations, 473
— — — (see also Hays, J. B.), 455
Korman, A. (see also Lee, G.), 407

λ (see also Bacteriophage), 1, 147, 163, 195, 277, 319, 473
lac promoter mutation p^r115 generates a new transcription initiation point, 551
— — mutations located downstream from the transcription start site, 537
Laskey, R. A. (see also Mills, A. D.), 561
Lee, G., Hannett, N. M., Korman, A. & Pero, J., Transcription of cloned DNA from *Bacillus subtilis* phage SP10. Requirement for hydroxymethyluracil-containing DNA by phage-modified RNA polymerase, 407
Lerman, L. S. (see also Robinson, B. H.), 19
Lifson, S. & Sander, C., Specific recognition in the tertiary structure of β-sheets of proteins, 627

McEwen, B. & Edelstein, S. J., Evidence for a mixed lattice in microtubules reassembled *in vitro*, 123
McKay, D. B. & Fried, M. G., Crystallization and preliminary X-ray diffraction data for the cyclic AMP receptor protein of *Escherichia coli*, 95
Mammalian mRNA, loss of the polyadenylate segment; selective cleavage of this sequence from polyribosomes, 439
Maquat, L. E. & Reznikoff, W. S., *lac* promoter mutation $P^{r}115$ generates a new transcription initiation point, 551
— — —., Thornton, K. & Reznikoff, W., *lac* promoter mutations located downstream from the transcription start site, 537
Marston, S. B. & Taylor, E. W., Comparison of the myosin and actomyosin ATPase mechanisms of the four types of vertebrate muscles, 573
Matthews, F. S. (see also Mercolino, T. J.), 557
Maturation, asymmetric, of a dimeric tRNA precursor, 349
Maurer, R., Meyer, B. J. & Ptashne, M., Gene regulation at the right operator (O_R) of bacteriophage λ. I. O_R3 and autogenous negative control by repressor, 147
Maurer, R. (see also Meyer, B. J.), 163
Melli, M. (see also Spinelli, G.), 111
Mercolino, T. J., Bellamy, H. D. & Matthews, F. S., A preliminary crystallographic study of β-glucuronidase from rat preputial gland, 557
Messenger RNA, fiber, incomplete splicing and deficient accumulation in monkey cells infected by human adenovirus 2, 221
—, histone, contained in high molecular weight RNA in sea urchin *Paracentrotus lividus*, 111
— RNA, mammalian, loss of the polyadenylate segment; selective cleavage of this sequence from polyribosomes, 439
— —, non-polyadenylated, in mouse Friend cells, coding potential, 61
Meyer, B. J. & Ptashne, M., Gene regulation at the right operator (O_R) of bacteriophage λ. III. λ repressor directly activates gene transcription, 195
— — —, Maurer, R. & Ptashne, M., Gene regulation at the right operator (O_R) of bacteriophage λ. II. O_R1, O_R2, and O_R3: their roles in mediating the effects of repressor and cro, 163
— — — (see also Maurer, R.), 147
Micelles, taurodeoxycholate, small-angle neutron scattering study of the association with porcine pancreatic colipase, 297
Microtubule assembly in the presence of guanosine diphosphate, equilibrium and kinetic analysis, 641
— —, role of co-operative interactions, microtubule-associated proteins and guanosine triphosphate; a model, 660
Microtubules reassembled *in vitro*, evidence for a mixed lattice, 123
Micro-video study of discontinuous growth of bacterial flagellar filaments *in vitro*, 265
Mills, A. D., Laskey, R. A., Black, P. & Robertis, E. M. de, An acidic protein which assembles nucleosomes *in vitro* is the most abundant protein in *Xenopus* oocyte nuclei, 561
Minty, A. J. & Gros, F., Coding potential of non-polyadenylated messenger RNA in mouse friend cells, 61
Mirzabekov, A. D. (see also Belyavsky, A. V.), 519
— — — (see also Shick, V. V.), 491
Mitochondrial DNA, human and mouse, analysis of sequence homology, 85
Monkey cells infected by human adenovirus type 2, incomplete splicing and deficient accumulation of the fiber mRNA, 221
Mouse Friend cells, coding potential of non-polyadenylated mRNA, 61
— and human mitochondrial DNA, analysis of their sequence homology, 85
MS2 (see also Bacteriophage), 243
Muscles, vertebrate, four types, a comparison of the myosin and actomyosin ATPase mechanisms, 573
Mutants, novel, of *Escherichia coli* that produce recombinogenic lesions in DNA; properties of recombinogenic λ phages grown on bacteria carrying *arl* mutations, 473
Mutation, azure, of bacteriophage MS2, 243
Mutations, novel, of *Escherichia coli* that produce recombinogenic lesions in DNA; identification and mapping of *arl* mutations, 455
Myosin and actomyosin ATPase mechanisms of the four types of vertebrate muscles, a comparison, 573

N-acetyl-D-neuraminic acid binding site location in wheat germ agglutinin, a crystallographic study at 2·8 Å resolution, 53

Neutron scattering study, small-angle, of the association between porcine pancreatic colipase and taurodeoxycholate micelles, 297
Nuclear magnetic resonance study of successively longer tRNA fragments with a common 5′ end; sequential folding of tRNA, 601
Nucleosome core particles, primary organization; sequential arrangement of histones along DNA, 491
— -assembling protein *in vitro* is the most abundant protein in *Xenopus* oocyte nuclei, 561
Nucleosomes containing all five histones and DNA 175 and 165 base-pairs long, primary organization, 519

O_R1, O_R2, and O_R3, their roles in mediating the effects of repressor and cro; gene regulation at the right operator of bacteriophage λ, 163
O_R3 and autogenous negative control by repressor; gene regulation at the right operator of bacteriophage λ, 147
Oocyte nuclei of *Xenopus*, an acidic protein which assembles nucleosomes *in vitro* is the most abundant protein, 561

P22 (see also Bacteriophage), 319
P^r115 *lac* promoter mutation generates a new transcription initiation point, 551
Paracentrotus lividus, high molecular weight RNA containing histone messenger, 111
Pero, J. (see also Lee, G.), 407
Phizicky, E. M. & Roberts, J. W., Kinetics of *recA* protein-directed inactivation of repressors of phage λ and phage P22, 319
Pkada, Y. (see also Iba, H.), 733
Polyadenylate segment, loss from mammalian mRNA; selective cleavage of this sequence from polyribosomes, 439
Porcine pancreatic colipase, small-angle neutron scattering study of the association with taurodeoxycholate micelles, 297
Precursor, dimeric, of tRNA: asymmetric maturation, 349
Preputial gland of the rat, a preliminary crystallographic study of β-glucuronidase, 557
Protein, acidic, which assembles nucleosomes *in vitro* is the most abundant protein in *Xenopus* oocyte nuclei, 561
—, outer surface, of bacteriophage lambda, 277
Proteins, specific recognition in the tertiary structure of β-sheets, 627
Ptashne, M. (see also Maurer, R.), 147
— — (see also Meyer, B. J.), 163, 190
Purple membrane, crystallization in three dimensions, 99

Rat preputial gland, a preliminary crystallographic study of β-glucuronidase, 557
Ray, B. K. & Apirion, D., Identification of RNA molecules which contain 5 S ribosomal RNA and transfer RNA in an RNAase E^- RNAase P^- double mutant strain of *Escherichia coli*, 329
Reassembled microtubules, evidence for a mixed lattice, 123
recA protein-directed inactivation kinetics of repressors of phage λ and phage P22, 319
Recognition, specific, in the tertiary structure of β-sheets of proteins, 627
Recombinogenic lesions produced in DNA by novel mutations of *Escherichia coli*; identification and mapping of *arl* mutations, 455
Recombinogenic lesions produced in DNA by novel mutations of *Escherichia coli*; properties of recombinogenic λ phages grown on bacteria carrying *arl* mutations, 473
Rementer, M. (see also Webster, R. E.), 393
Replication of bacteriophage f1: a complex containing gene *II* protein in gene *V* mutant-infected bacteria, 393
Repressors of phage λ and phage P22, kinetics of *recA* protein-directed inactivation, 319
Reznikoff, W. S. (see also Maquat, L. E.), 551
RNA, high molecular weight, containing histone messenger in the sea urchin *Paracentrotus lividus*, 111
— molecules containing 5 S ribosomal RNA and tRNA in an RNAase E^- RNAase P^- double mutant strain of *Escherichia coli*, 329
— polymerase, phage-modified, requires hydroxymethyluracil-containing DNA for transcription; transcription of cloned DNA from *Bacillus subtilis* phage SP01, 407

Ribosomal 5 S RNA and tRNA are contained in RNA molecules in an RNAase E⁻ RNAase P⁻ double mutant strain of *Escherichia coli*, 329
RNAase E⁻ RNAase P⁻ double mutant strain of *Escherichia coli*, identification of RNA molecules which contain 5 S ribosomal RNA and tRNA, 329
Robertis, E. M. de (see also Mills, A. D.), 561
Roberts, J. W. (see also Phizicky, E. M.), 319
Robillard, G.T. (see also Boyle, J.), 601
Robinson, B. H., Lerman, L. S., Beth, A. H., Frisch, H. L., Dalton, L. R. & Auer, C., Analysis of double-helix motions with spin-labeled probes: binding geometry and the limits of torsional elasticity, 19

Saccharomyces cerevisiae, a *cis*-dominant regulatory mutation linked to the *argB-argC* gene cluster, 691
Sander, C. (see also Lifson, S.), 627
Schizosaccharomyces pombe, the antisuppressor strain *sin1* lacks the modification isopentenyladenosine in tRNA, 207
Scholla, C. A. (see also Guthrie, C.), 349
Sea urchin *Paracentrotus lividus*, high molecular weight RNA containing histone messenger, 111
Séméríva, M. (see also Charles, M.), 297
Sequence, DNA, of a T4 tRNA gene cluster, 377
— homology between human and mouse mitochondrial DNA, an analysis, 85
Sequential folding of transfer RNA; n.m.r. study of successively longer tRNA fragments with a common 5′ end, 601
Serine enzyme *Streptomyces griseus* protease A and the bacterial-produced inhibitor chymostatin, structure of the complex formed between them, 45
Shick, V. V., Belyavsky, A. V., Bavykin, S. G. & Mirzabekov, A. D., Primary organization of the nucleosome core particles. Sequential arrangement of histones along DNA, 491
Shotton, D. (see also Henderson, R.), 99
Sielecki, A. R. (see also James, M. N. G.), 423
sin1 strain of *Schizosaccharomyces pombe* lacks the modification isopentenyladenosine in tRNA, 207
Soll, L. (see also Knowlton, R. G.), 705
Specificity, dual, of su^+7 tRNA. Evidence for translational discrimination, 705
Spinelli, G., Melli, M., Arnold, E., Casano, C., Gianguzza, F. & Ciaccio, M., High molecular weight RNA containing histone messenger in the sea urchin *Paracentrotus lividus*, 111
Spin-labeled compound preparation, 41
— — probes and the analysis of double helix motions: binding geometry and the limit of torsional elasticity, 19
Splicing, incomplete, and deficient accumulation of the fiber mRNA in monkey cells infected by human adenovirus type 2, 221
SP01 (see also Bacteriophage), 407
Streptomyces griseus protease A, structure of the complex formed with the bacterial-produced inhibitor chymostatin, 45
— — — B and tripeptide chloromethyl ketone inhibitors, crystal structure studies and inhibition kinetics, 423
Structure of the complex between the bacterial-produced inhibitor chymostatin and the serine enzyme *Streptomyces griseus* protease A, 45
— — — — between wheat germ agglutinin and *N*-acetyl-D-neuraminic acid, 53
Structure of the cyclic AMP receptor protein of *Escherichia coli*, 95
— — — nucleosome core particles; sequential arrangement of histones along DNA, 491
— — nucleosomes containing all five histones and DNA 175 and 165 base-pairs long, 519
— — reassembled microtubules, 123
— ,tertiary, of β-sheets of proteins, specific recognition, 627
su^+7 tRNA dual specificity; evidence for translational discrimination, 705
— —, discrimination between aminoacyl groups by elongation factor Tu, 721
Surface protein, outer, of bacteriophage lambda, 277
Syvanen, M., Tn903 induces inverted duplications in the chromosome of bacteriophage lambda, 1

T4 (see also Bacteriophage), 377, 679
Tail fiber function in triggering baseplate expansion of bacteriophage T4, 679
Taurodeoxycholate micelles, small-angle neutron scattering study of the association with porcine pancreatic colipase, 297

Taylor, E. W. (see also Marston, S. B.), 573
Thornton, K. (see also Maquat, L. E.), 537
Tn903 induces inverted duplications in the chromosome of bacteriophage lambda, 1
Torsional elasticity limit and binding geometry; analysis of double-helix motions with spin-labeled probes, 19
Transcription initiation point, new, is generated by the *lac* promoter mutation p^r115, 551
— of cloned DNA from *Bacillus subtilis* phage SPO1. Requirement for hydroxymethyluracil-containing DNA by phage-modified RNA polymerase, 407
— start site, *lac* promoter mutations located downstream, 537
Transfer RNA from the antisuppressor strain *sin1* of *Schizosaccharomyces pombe*, the modification isopentenyladenosine is missing, 207
— — gene cluster, T4, determination of DNA sequence, 377
— — precursor, dimeric, asymmetric maturation, 349
— — and 5 S ribosomal RNA are contained in RNA molecules in an RNAase E^- RNAase P^- double mutant strain of *Escherichia coli*, 329
— — sequential folding; n.m.r. study of successively longer tRNA fragments with a common 5′ end, 601
— —, su^+7, discrimination between aminoacyl groups by elongation factor Tu, 721
— — ,su^+7, dual specificity; evidence for translational discrimination, 705
Translational discrimination evidence; dual specificity of su^+7 tRNA, 705
Tripeptide chloromethyl ketone inhibitors, crystal structure studies and inhibition kinetics with *Streptomyces griseus* protease B, 423
Tsugita, R. (see also Imber, R.), 277
Tu discriminates between aminoacyl groups on su^+7 tRNA, 721

Van Montagu, M. (see also Iserentant, D.), 243
Vertebrate muscles, a comparison of the myosin and actomyosin ATPase mechanisms of the four types, 573
Virus (see also Adenovirus type 2), 221
Vögeli, G. (see also Janner, F.), 207

Webster, R. E. & Rementer, M., Replication of bacteriophage f1: a complex containing gene *II* protein in gene *V* mutant-infected bacteria, 393
Weisenberg, R. C., Role of co-operative interactions, microtubule-associated proteins and guanosine triphosphate in microtubule assembly: a model (*Appendix* to Zackroff, R. V. *et al.*, 641), 660
— — (see also Zackroff, R. V.), 641
Wheat germ agglutinin, location of the *N*-acetyl-D-neuraminic acid binding site; crystallographic study at 2·8 Å resolution, 53
Wright, C. S., Location of the *N*–acetyl-D-neuraminic acid binding site in wheat germ agglutin. A crystallographic study at 2·8 Å resolution, 53
Wurtz, M. (see also Imber, R.), 277

Xenopus oocyte nuclei, an acidic protein which assembles nucleosomes *in vitro* is the most abundant protein, 561
X-ray diffraction data, preliminary, and crystallization for the cyclic AMP receptor protein of *Escherichia coli*, 95

Yarus, M. (see also Knowlton, R. G.), 705, 721

Zackroff, R. V., Weisenberg, R. C. & Deery, J., Equilibrium and kinetic analysis of microtubule assembly in the presence of guanosine diphosphate, 641